ANBCA 2

Analytical Biochemistry

An International Journal

UNIVERSITY OF HAWAII LIBRARY
LIBRARY USE ONLY

EDITOR: *Alvin Nason*

ADVISORY BOARD:

Gary Ackers
Norman G. Anderson
Eraldo Antonini
Gilbert Ashwell
Samuel P. Bessman
Simon Black
Robert M. Bock
Robert L. Bowman
Ludwig Brand
A. E. Braunstein
Melvin Calvin
Andreas Chrambach
Sidney Colowick
Frank Eisenberg, Jr.
Jacques R. Fresco
Victor Ginsburg
I. C. Gunsalus
Harlyn O. Halvorson
William Z. Hassid
Osamu Hayaishi
Edward C. Heath
C. H. W. Hirs
Evan C. Horning
Rollin D. Hotchkiss
Walter L. Hughes
Ephraim Katchalski
Irving Klotz
Luis F. Leloir
Olov Lindberg
Feodor Lynen
Vincent Massey
Manfred M. Mayer
Bruce Merrifield
Marshall Nirenberg
Kazuo Okunuki
Alexander Rich
Gerhard Schmidt
Michael Sela
Herbert A. Sober
Sol Spiegelman
Paul K. Stumpf
Charles A. Thomas, Jr.
Sidney Udenfriend
Bert L. Vallee
Donald D. Van Slyke
Samuel B. Weiss
Theodor Wieland
George Wolf
W. A. Wood
Lemuel D. Wright
David A. Yphantis

VOLUME 40
1971

ACADEMIC PRESS
New York and London

Copyright ©, 1971, by Academic Press, Inc.
ALL RIGHTS RESERVED
No part of this volume may be reproduced in any form, by photostat, microfilm, by retrieval system, or any other means, without written permission from the Publishers.

Made in the United States of America

CONTENTS OF VOLUME 40

Number 1, March 1971

G. J. E. Balharry and D. J. D. Nicholas. New Assay for ATP-Sulphurylase Using the Luciferin-Luciferase Method . . . 1

Tao Wong. Studies on Creatine Determination by α-Naphthol-Diacetyl Reaction 18

Kenneth J. Stevenson. Use of Visual Dye Markers on High-Voltage Paper Ionophoresis 29

C.-P. Kedenburg. A Lithium Buffer System for Accelerated Single-Column Amino Acid Analysis in Physiological Fluids . 35

E. G. Richards and R. Lecanidou. Quantitative Aspects of the Electrophoresis of RNA in Polyacrylamide Gels 43

Walton H. Marsh and Patrick J. Fitzgerald. Pancreas Acinar Cell Differentiation. IV. Assay of Nanogram Levels of Chymotrypsin by Radioactively Labeled Synthetic Substrate . . 72

L. H. Krull, D. E. Gibbs, and Mendel Friedman. 2-Vinylquinoline, a Reagent to Determine Protein Sulfhydryl Groups Spectrophotometrically 80

Frederic L. Hoch, Rosemary A. Kuras, and John D. Jones. Iodine Analysis of Biological Samples by Neutron Activation of ^{127}I, with Scintillation Counting of Cerenkov Radiation . 86

D. Rodbard and A. Chrambach. Estimation of Molecular Radius, Free Mobility, and Valence Using Polyacrylamide Gel Electrophoresis 95

D. Rodbard, G. Kapadia, and A. Chrambach. Pore Gradient Electrophoresis 135

J. Lunney, A. Chrambach, and D. Rodbard. Factors Affecting Resolution, Band Width, Number of Theoretical Plates, and Apparent Diffusion Coefficient in Polyacrylamide Gel Electrophoresis 158

John P. Mallon and Colin Dalton. Automated Enzymic Method for the Determination of Glycerol 174

N. B. Cucakovich. Determination of Tween 80 in Tissue Culture Media, Vaccines, and Related Products 183

J. Korf, S. Ottema, and I. van der Veen. Fluorometric Determination of Homovanillic Acid in Biological Material after Isolation of Sephadex G-10 187

D. L. Alkon, Arthur S. Goldberg, Jack Peter Green, Patrizia Levi, and Karen Liao. Fluorophors Formed from Imidazoles 192

Naoyuki Taniguchi. Quantitative Microdetermination of Total Cysteine plus Cystine in Proteins 200

Dolores Bobb and B. H. J. Hofstee. Gel Isoelectric Focusing

for Following the Successive Carbamylations of Amino Groups in Chymotrypsinogen A 209
Elisha Haas, Yehudit Elkana, and Richard G. Kulka. A Sensitive Fluorometric Assay for α-Chymotrypsin 218

Short Communications

Per Hellung-Larsen and Sune Frederiksen. Correlation Between Separation of Low Molecular Weight RNA on MAK Columns and Polyacrylamide Gels 227
B. G. Johansson and J. Stenflo. Polyacrylamide Slab Electrophoresis Followed by Electrophoresis into Antibody-Containing Agarose Gel 232
P. W. Atkinson, W. V. Brown, and A. R. Gilby. Gas Chromatographic Analysis of *N*-Acetyldopamine and Its Potential Metabolic Precursors 236

Number 2, April 1971

Norman R. Lazarus, Raul A. Gutman, and Lillian Recant. A Method for Electrophoretic Characterization on Polyacrylamide Gel of Circulating Insulin Immunoreactive Substances 241
M. Popescu, L. H. Lazarus, and N. Goldblum. Simplified Adaptor for Electroelution 247
Charles P. Emerson, Jr. and Tom Humphreys. A Simple and Sensitive Method for Quantitative Measurement of Cellular RNA Synthesis 254
Nino L. Incardona, Harvey Notarius, and J. Bert Flanegan. Measurement of Temperature within the Sample Cell during Sedimentation Velocity Experiments 267
Nils-Georg Asp. Improved Method for the Assay of Phenylglycosidase Activity with a 4-Aminoantipyrine Reagent . . 281
James S. Whitehead, Ernest Kay, Jow Y. Lew, and Leland M. Shannon. A Preparative Column Electrophoresis Apparatus Using Sephadex G-25 287
G. R. Finlayson and A. Chrambach. Isoelectric Focusing in Polyacrylamide Gel and Its Preparative Application . . . 292
David Cooney, Ruth Davis, and Glen Van Atta. A Spectrophotometric Method for the Simultaneous Measurement of L-Glutamine and L-Asparagine in Biological Materials . . 312
Fritz Heinz and Wolfgang Prosch. Estimation of the Molecular Weight of Polypeptide Chains by Thin-Layer Gel-Chromatography in 6 *M* Guanidine Hydrochloride 327
Bernardo S. Vanderheiden. Preparation of Labeled Nucleotides.

III. ^{14}C-Labeled Adenosine, Guanosine, and Inosine Phosphates, and ^{32}P-Labeled Inosine Monophosphate 331
Sigmund F. Zakrzewski and Jaakko J. Himberg. Preparation of Tritiated Dihydrofolate and Its Use in the Assay of Dihydrofolate Reductase 336
James R. Florini, Rosaria P. Brivio, and Barbara-Anne M. Battelle. Isoelectric Focusing of Myosin in Dilute Polyacrylamide Gels 345
R. Libert and R. de Hertogh. 2,4-Dinitrophenylhydrazones of 17-Ketosteroids. Description of Some Physicochemical Properties and Their Practical Application in Quantitative Measurements 351
G. L. Nelsestuen and S. Kirkwood. A Simple, Economical Synthesis of Uridine Diphospho-*N*-acetylgalactosamine-6-^{3}H . . 359
Michael T. Ryan and Nabil S. Hanna. Investigation of Equilibrium Ultrafiltration as a Means of Measuring Steroid-Protein Binding Parameters 364
Joel K. Weltman. A Simplified Model for Analysis of Conversion of Methionine to Cysteine *in Vivo*. 380
C. L. Long, Y. Mashima, and F. Gump. A Simple Method for Measurement of Specific Activity of C^{14}-Lactate 386
Douglas B. Swartz and George S. Wilson. Small-Volume Coulometric Redoxostat 392
Charles W. Lees. Procedure for the Base Composition of RNA Especially Suited for Samples Containing Relatively Large Amounts of DNA and Protein 401
Abraham Saifer. Comparative Study of Various Extraction Methods for the Quantitative Determination of Free Amino Acids from Brain Tissue 412
Syed N. Alam and Ross H. Hall. Separation of the Δ^2 and Δ^3 Isomers of N^6-(Δ^2-Isopentenyl)-adenosine by Gas Chromatography and Characterization by Mass Spectroscopy . 424
Bohdan Bakay. Detection of Radioactive Components in Polyacrylamide Gel Disc Electropherograms by Automated Mechanical Fractionation 429
Jeremy S. D. Winter and Derek R. Grant. A Rapid and Sensitive Assay for Plasma Testosterone in Adults and Children 440
Paul Carter. Spectrophotometric Determination of Serum Iron at the Submicrogram Level with a New Reagent (Ferrozine) 450
Kin-Ping Wong, Robert Roxby, and Charles Tanford. Criteria of Purity for Guanidine Hydrochloride 459
William Niedermeier. Gas Chromatography of Neutral and Amino Sugars in Glycoproteins 465

Short Communications

R. V. Josephson, S. K. Maheswaran, C. V. Morr, R. Jenness, and R. K. Lindorfer. Effect of Urea on p*I*'s of Ampholytes and Casein in Isoelectric Focusing 476

A. R. Henderson. Longitudinal Slicing of Polyacrylamide Slab Gels: A Simple Technique 482

Ashima Chakravarti and John W. Thanassi. Liquid Scintillation Counting of C^{14} and H^3 Samples on Filter Paper . 484

Shingo Marumo, Hiroyuki Hattori, and Hiroshi Abe. Chromatography of a New Natural Auxin, 4-Chloroindolyl-3-acetic Acid and Related Chloro Derivatives . 488

Paul T. Englund, Hamilton O. Smith, and Earl Sandbek. A Density Gradient Fractionator 490

Author Index 494

Subject Index 496

r man from quality control is real mean.

because he knows that the tion of Boehringer Mann- iochemicals is based on the level of quality control.
n because, with this knowl- the total enzyme system, he ands the needs of basic bio- al research.
is why he concentrates his on the biochemical require- for the total enzyme system e, coenzyme and substrate). he case of enzymes, stating inants in relation to the spe- tion of the enzyme, he insists optimal conditions for the ation of the contaminant be

coenzymes and substrates; n insists that they are bio- ally assayed, rather than by hemical and/or physical ss.
all our Quality Control Man but he calls it awareness.... act BMC for the Biochemi- talogue containing over 250 mical assay and specification . This catalog includes 30 new cts recently introduced, that just meet your research re- ments.

DIAGNOSTICS/BIOCHEMICALS

bmc

NGER MANNHEIM CORPORATION
44TH STREET, NEW YORK, N. Y. 10017

If it's on the chromatogram it's on the recording. All of it.

EXACTLY.

Our chromatogram-to-data converter: the Schoeffel SD 3000. You can use it for quantitative and qualitative analyses of TLC plates, gels, paper strips, electrophoretograms, films, radiogram plates, etc., etc. Nothing is more accurate, nothing more sensitive, nothing more versatile.

More SD 3000's are in use than all other research grade chromatogram analysis systems combined.

The SD 3000 is the only totally intercepting, double-beam, ratio recording chromatogram analyzer capable of true quantitation. It operates in transmission (% T, linear O.D.) or reflection (fluorescence, absorption, fluorescence quenching) modes. It has a high intensity light source, monochromators, matched photomultiplier detectors, density computer, integrating recorder. Just about any accessory you might need. It provides continuous spectral scanning over a 200 nm to 700 nm (UV-VIS) range. The SD 3000 permits rapid, automatic, non-destructive analyses of sub-nanogram samples (or preparative quantities) with resolution and reliability hardly approached in any other system.

Care to send us a plate for analysis? A prompt demonstration in your laboratory? Literature? No problem. Just write or 'phone.

Schoeffel Instrument Corp., 24 Booker Street, Westwood, N.J. (201) 664-7263. In Europe: Schoeffel Instrument GMBH, 2351 Trappenkamp, C Strasse 2, Deutschland.



SCHOEFFEL
LIGHT AND LIGHT MEASUREMENT

THE UN-NEW SPECTROPHOTOMETER



gilford®

FLEXIBILITY TO MEET YOUR GROWING NEEDS

FOR GEL SCANNING

- Proven Performance — Hundreds in use
- High Resolution — Interchangeable micro-apertures
- Easy and Fast to Operate and Install



FOR RAPID SAMPLING

- New Sample Return Feature
- Automatic Print-Out
- Up to 250 Samples per Hour



FOR ACCURATE TEMPERATURE CONTROL

- Totally ELECTRONIC Heating AND Cooling
- Precision Thermostatting and Controlled Heating Rates
- Rapid Cooling Rates — for Renaturation and Recycling

See these and other new developments at FASEB BOOTHS G and F, 31 and 32, or write:

GILFORD INSTRUMENT LABORATORIES, INC., OBERLIN, OHIO 44074

GILFORD EUROPE, S.A. • 74 RUE M. S. DEFRESNE • 94 VITRY-SUR-SEINE, FRANCE

For precision and speed, Model KLA-3B Amino Acid Analyzer passes the acid test.

The Perkin-Elmer Model KLA-3B features the exclusive Ligand Exchange System for the most rapid and precise separation of simple or complex amino acid samples. Finished accelerated chromatograms provide *unexcelled peak-to-valley resolution of such difficult components as threonine and serine.*

Automated Operation. Provides 24 and more hours of continuous unattended operation with 24 separate samples automatically introduced and analyzed. *Offers assembly line convenience and lower per sample cost.*

Continuous Flow Pulse-Free Pumping. Three stainless steel minimum flow pumps deliver constant flow streams independent of system back pressures or eluent viscosities. These pumps are unique among precision chromatographic pumps, *providing smoother baseline and greater precision.*

Automatic Column Regeneration. All regeneration cycles are completely overlapped with the sample analysis cycle. No waiting after each sample for regeneration and flushing thereby *providing savings of time and money.*

Triple Wavelength Analysis. Covers a wider dynamic range of component concentration at one time. Novel single-source triple detector photometer furnishes signals to the recorder from three different wavelengths. Monitors amino and imino acids at 440 $m\mu$, 570 $m\mu$ and 640 $m\mu$ for *complete spectrum coverage to give more useful data per sample.*

For detailed information on the Perkin-Elmer Model KLA-3B write to: Instrument Division, Perkin-Elmer Corporation, XXX Main Avenue, Norwalk, Conn. 06852.



PERKIN-ELMER

Acids—^{14}C Labeled
warz/Mann
ngeburg, New York 10962
4) 359-2700
ADVERTISEMENT THIS PAGE)

Acids—^{3}H Labeled
warz/Mann
ngeburg, New York 10962
4) 359-2700
ADVERTISEMENT THIS PAGE)

Acids—
d ^{35}S Labeled
warz/Mann
ngeburg, New York 10962
4) 359-2700
TE FOR INFORMATION)

Acids—
stituted Protein
lysates (^{14}C, ^{3}H)
warz/Mann
ngeburg, New York 10962
4) 359-2700
TE FOR INFORMATION)

Acids—
l Contract Orders
warz/Mann
ngeburg, New York 10962
4) 359-2700
TE FOR INFORMATION)

Acids—t-Boc
warz/Mann
supply virtually every
ting material needed
peptide synthesis:
Boc-amino acids, 22 ^{14}C
^{3}H labeled Boc-amino
s, 12 Boc-amino acid
n esters, etc. Write or
for brochure.
ngeburg, New York
62 (914) 359-2700

Acids—Unlabeled
warz/Mann
ngeburg, New York 10962
4) 359-2700
TE FOR INFORMATION)

Acids

^{14}C Amino Acids from Schwarz/Mann

DIVISION OF BECTON, DICKINSON AND COMPANY

Examples of the unusually high specific activities available from Schwarz/Mann:

L-[^{14}C] Alanine	159mc/mM	L-[^{14}C] Lysine	320mc/mM
L-[^{14}C] Arginine	320mc/mM	L-[^{14}C] Methionine	255mc/mM
L-[^{14}C] Aspartic Acid	212mc/mM	L-[^{14}C] Phenylalanine	455mc/mM
[^{14}C] Glycine	106mc/mM	L-[^{14}C] Threonine	212mc/mM
L-[^{14}C] Leucine	320mc/mM	L-[^{14}C] Tyrosine	455mc/mM

Further re our high specific activities: ALL of our ^{14}C amino acids are higher than 50mc/mA. About purity. Every lot of phenylalanine and tyrosine is repurified every month. Every lot of everything else: reassayed every ten weeks and repurified if necessary. And *all* assay and reassay data goes onto the Product Analysis Report that you receive. About availability: these compounds are available *from stock*. For the complete story on our complete labeled amino acid line call (914) 359-2700 or write: Schwarz/Mann, Orangeburg, New York 10962. Division of Becton, Dickinson and Company.

(914) 359-2700

(SEE OUR OTHER ADVERTISEMENT AND LISTINGS ON THIS PAGE.)

^{3}H AMINO ACIDS FROM SCHWARZ/MANN

Examples of the unusually high specific activities available from Schwarz/Mann:

Catalytic

L-[^{3}H] Alanine	6c/mM
L-[^{3}H] Arginine	10c/mM
L-[^{3}H] Histidine	8c/mM
L-[^{3}H] Lysine	6c/mM
L-[^{3}H] Phenylalanine	7c/mM
L-[^{3}H] Valine	8c/mM

Specifically Labeled

L-[4^{3}H] Arginine	20c/mM
L-[2,3^{3}H] Aspartic Acid	15c/mM
L-[3,4^{3}H] Glutamic Acid	15c/mM
L-[4,5^{3}H] Leucine	50c/mM
L-[4,5^{3}H] Lysine	20c/mM
L-[5^{3}H] Proline	15c/mM
L-[2,3^{3}H] Valine	15c/mM

These high specific activity ^{3}H-labeled amino acids are reassayed every ten weeks—as is *every* radiochemical we make—and repurified when necessary. All relevant analytical data gets forwarded to you. For detailed information call (914) 359-2700 or write: Schwarz/Mann, Orangeburg, New York 10962. Division of Becton, Dickinson and Company.

(SEE OUR OTHER ADVERTISEMENT AND LISTINGS ON THIS PAGE.)

For the rest of the Schwarz/Mann amino acid story write or call Schwarz/Mann Orangeburg, N.Y. 10962 (914) 359-2700



Another First from Sigma

COMPLETE OFFERING

OF ENZYMES AND SUBSTRATES FOR

"PENTOSE SHUNT"

For the first time, *all* enzymes and substrates for "Pentose Shunt" are available. Some compounds have been offered by Sigma before. The following preparations, however, are offered here for the first time anywhere.

Unit Definition for Enzymes: One Unit will convert 1 μMole of Substrate per minute at 25°C. Data sheet will accompany each compound.

TRANSKETOLASE

T9006 Type III Crystalline From Baker's Yeast Ammonium Sulfate Suspension

Activity: approx. 15 Units/mg protein

Impurities determined: ADH approx. 1% Ru5P3-Epimerase <0.1% Phosphoriboisomerase <0.1% Transaldolase <0.1%

5 units	$13.20	50 units	$100.00
10 units	22.00	100 units	180.00

T9381 Type IV Crystalline From Baker's Yeast Ammonium Sulfate Suspension

Activity: approx. 20 Units/mg protein

Impurities determined: ADH <0.1% Ru5P3-Epimerase <0.1% Phosphoribolsomerase <0.1% Transaldolase <0.1%

5 units	$18.00	50 units	$135.00
10 units	30.00	100 units	250.00

D-ERYTHROSE-4-PHOSPHATE (E4P)

E0377 Grade II Semi-purified Sodium Salt
Purity: approx. 70% Mol. Wt.: 244

It replaces the old Dimethylacetal Dicyclohexyl-ammonium Salt (Mol. Wt. 462) and does not require further hydrolysis.

2 mg	$ 5.80	25 mg	$ 31.50
5 mg	10.00	100 mg	100.00
		250 mg	$200.00

To be released soon: Higher purity Phosphoribo-somerase, Ribulose-5-Phosphate-3-Epimerase, Xy-lulose-5-Phosphate, Sedoheptulose-7-Phosphate.

TRANSALDOLASE

T9506 Grade III Crystalline From Torula Yeast Ammonium Sulfate Suspension
Activity: 15–30 units/mg protein

5 units	$9.00	25 units	$30.00
		100 units	$80.00

RIBULOSE-5-PHOSPHATE-3-EPIMERASE

R0626 Type II From Baker's Yeast
Ammonium Sulfate Suspension
Activity: approx. 5 units/mg protein
This is a rather impure preparation at present, but it seems to be quite adequate for many studies.

2 units	$12.00	5 units	$24.00
		20 units	$80.00

XYLULOSE-5-PHOSPHATE (Xu5P)

X3125 Crude Sodium Salt
Purity: Approx. 25% (balance mostly R5P).
Although crude, it seems to be quite adequate for use as substrate for Transketolase.

5 mg	$ 3.80	100 mg	$ 36.00
25 mg	12.90	500 mg	120.00
		1 g	$200.00

SEDOHEPTULOSE-7-PHOSPHATE (S7P)

S0878 Grade II Barium Salt
Purity: approx. 85%.
Impurities determined: G6P approx. 2% Triosephosphates approx. 2% F6P <1%

2 mg	$ 5.80	25 mg	$ 31.50
5 mg	10.00	100 mg	100.00
		250 mg	$200.00

Please consult our latest catalog for details of the following additional reagents for the "Pentose Shunt":

GLUCOSE-6-PHOSPHATE DEHYDROGENASE
6-PHOSPHOGLUCONIC DEHYDROGENASE
PHOSPHOGLUCOSE ISOMERASE
PHOSPHORIBOSOMERASE
FRUCTOSE-6-PHOSPHATE (F6P)
GLUCOSE-6-PHOSPHATE (G6P)
GLYCERALDEHYDE-3-PHOSPHORIC ACID (G3AP)
6-PHOSPHOGLUCONIC ACID (6PG)
RIBOSE-5-PHOSPHATE (R5P)
RIBULOSE-5-PHOSPHATE (Ru5P)

It's a pleasure doing business with Sigma

ORDER *DIRECT*—TELEPHONE *COLLECT* from ANYWHERE in the WORLD

Day, *Station to Station*, 314/771-5750 • Night, *Person to Person*, Dan Broida, 314/993-6418

TWX (Teletype) Day or Night: COLLECT-910-761-0593

TELEGRAM: SIGMACHEM, St. Louis, Missouri



The Research Laboratories of

SIGMA® CHEMICAL COMPANY

MAILING ADDRESS: P. O. BOX 14508, ST. LOUIS, MO., 63178, U.S.A.

MANUFACTURERS OF THE FINEST BIOCHEMICALS AVAILABLE

Distributed in the United Kingdom through
SIGMA LONDON Chem. Co. Ltd. • 12, Lettice St., London, S.W. 6, England • Phone 01-736-5823 (Reverse Charges)